

Contents lists available at [ScienceDirect](http://www.sciencedirect.com)

Urological Science

journal homepage: www.urol-sci.com

Corrigendum to “Results of a randomized, double-blind, placebo-controlled study of mirabegron in a Taiwanese population with overactive bladder and comparison with other clinical trials” [Urol Sci (2015) 41–48]



Hann-Chorng Kuo ^{a,*}, Ho-Hsiung Lin ^b, Hong-Jeng Yu ^c, Chen-Li Cheng ^d,
Man-Jung Hung ^e, Alex Tong-Long Lin ^f, The Taiwan Mirabegron Study Team^g

^a Department of Urology, Buddhist Tzu Chi General Hospital and Tzu Chi University, Hualien, Taiwan

^b Department of Obstetrics and Gynecology, National Taiwan University Hospital, Taipei, Taiwan

^c Department of Urology, National Taiwan University Hospital, Taipei, Taiwan

^d Department of Urology, Taichung Veterans General Hospital, Taichung, Taiwan

^e Department of Obstetrics and Gynecology, Taichung Veterans General Hospital, Taichung, Taiwan

^f Department of Urology, Taipei Veterans General Hospital, Taipei, Taiwan

The conflicts of interest statement at the end of this article is hereby corrected to:

This study was an analysis of the Taiwanese data in the Asian Mirabegron Trial. Although Astellas sponsored the Asian Mirabegron Trial, Astellas did not have any input on any aspect of the study. We declare that we have no conflicts of interest with Astellas in relation to this work.

DOI of original article: <http://dx.doi.org/10.1016/j.urols.2014.12.010>.

* Corresponding author. Department of Urology, Buddhist Tzu Chi General Hospital, Number 707, Section 3, Chung-Yang Road, Hualien 970, Taiwan.

E-mail address: hck@tzuchi.com.tw (H.-C. Kuo).

^g **The Taiwan Mirabegron Study Team:** Hann-Chorng Kuo, Department of Urology, Buddhist Tzu Chi General Hospital and Tzu Chi University, Hualien, Taiwan; Ho-Hsiung Lin, Department of Obstetrics and Gynecology, National Taiwan University Hospital, Taipei, Taiwan; Hong-Jeng Yu, Department of Urology, National Taiwan University Hospital, Taipei, Taiwan; Chen-Li Cheng, Department of Urology, Taichung Veterans General Hospital, Taichung, Taiwan; Man-Jung Hung, Department of Obstetrics and Gynecology, Taichung Veterans General Hospital, Taichung, Taiwan; Alex Tong-Long Lin, Department of Urology, Taipei Veterans General Hospital, Taipei, Taiwan; Chao-Hsiang Chang, Department of Urology, China Medical University Hospital, Taichung, Taiwan; Yao-Chi Chuang, Department of Urology, Kaohsiung Chang Gung Memorial Hospital, Kaohsiung, Taiwan; Tai-Lung Cha, Department of Urology, Tri-Service General Hospital, Taipei, Taiwan; Gin-Den Chen, Department of Obstetrics and Gynecology, Chung Shan Medical University Hospital, Taichung, Taiwan; Chih-Shou Chen, Department of Urology, Chiayi Chang Gung Memorial, Chiayi, Taiwan; Ming-Ping Wu, Department of Obstetrics and Gynecology, Chi Mei Foundation Hospital, Tainan, Taiwan; Tong-Lin Wu, Department of Urology, Kaohsiung Veterans General Hospital, Kaohsiung, Taiwan; Ken-Jen Yu, Department of Obstetrics and Gynecology, Kaohsiung Veterans General Hospital, Kaohsiung, Taiwan; Shih-Tsung Huang, Department of Urology, Linkou Chang Gung Memorial Hospital, Taoyuan, Taiwan.

<http://dx.doi.org/10.1016/j.urols.2015.06.290>

1879-5226/Copyright © 2015, Taiwan Urological Association. Published by Elsevier Taiwan LLC. Open access under [CC BY-NC-ND license](http://creativecommons.org/licenses/by-nc-nd/4.0/).